

# Sub- and super-fidelity as bounds for quantum fidelity

J. A. Miszczak, Z. Puchała, P. Horodecki,  
A. Uhlmann, K. Życzkowski

October 28, 2008

## Introduction

- Quantum states
- Properties of fidelity
- Analysis of fidelity

## Bounds for fidelity

- Fidelity, sub- and super-fidelity
- Properties of sub- and super-fidelity
- Comparison of different bounds
- Super-fidelity and trace distance
- Measuring and calculating super-fidelity

## Conclusions

## References

## Quantum states and fidelity

Quantum state is an operator  $\rho : \mathcal{H}_N \rightarrow \mathcal{H}_N$ , which is positive semi-definite ( $\rho \geq 0$ ) and normalised ( $\text{tr} \rho = 1$ ). We denote by  $\Omega_N \subset \mathcal{M}_N$  the space of density matrices on  $\mathbb{C}^N$ .

We define the fidelity between two states as

$$F(\rho_1, \rho_2) = (\text{tr} |\sqrt{\rho_1} \sqrt{\rho_2}|)^2 = \|\rho_1^{1/2} \rho_2^{1/2}\|_1^2,$$

where  $\|\cdot\|_1$  is a trace norm, *i.e.*  $\|A\|_1 = \text{tr}|A| = \sum_{i=1}^N \sigma_i(A)$ .

In the case of two pure states  $\rho_1 = |\phi\rangle\langle\phi|$ ,  $\rho_2 = |\psi\rangle\langle\psi|$  we have  $F(\rho_1, \rho_2) = |\langle\psi|\phi\rangle|^2$ .

# Properties of fidelity

Fidelity has few nice properties

- ▶ **Bounds:**  $0 \leq F(\rho_1, \rho_2) \leq 1$ . Furthermore  $F(\rho_1, \rho_2) = 1$  iff  $\rho_1 = \rho_2$ , while  $F(\rho_1, \rho_2) = 0$  iff  $\text{supp}(\rho_1) \perp \text{supp}(\rho_2)$ .
- ▶ **Symmetry:**  $F(\rho_1, \rho_2) = F(\rho_2, \rho_1)$ .
- ▶ **Unitary invariance:**  $F(\rho_1, \rho_2) = F(U\rho_1U^\dagger, U\rho_2U^\dagger)$ , for any unitary operator  $U$ .
- ▶ **Concavity:**  
 $F(\rho, a\rho_1 + (1-a)\rho_2) \geq aF(\rho, \rho_1) + (1-a)F(\rho, \rho_2)$ , for  $a \in [0, 1]$ .
- ▶ **Multiplicativity:**  $F(\rho_1 \otimes \rho_2, \rho_3 \otimes \rho_4) = F(\rho_1, \rho_3)F(\rho_2, \rho_4)$ .
- ▶ **Joint concavity:**  $\sqrt{F}(a\rho_1 + (1-a)\rho_2, a\rho'_1 + (1-a)\rho'_2) \geq a\sqrt{F}(\rho_1, \rho'_1) + (1-a)\sqrt{F}(\rho_2, \rho'_2)$ , for  $a \in [0, 1]$ .

# Classical counterpart

Fidelity between two diagonal operators is equal to the *Bhattacharyya* coefficient  $B$  for their eigenvalues.

$$\sqrt{F(\text{diag}(\rho_1), \text{diag}(\rho_2))} = B(p, q) = \sum_{i=1}^n \sqrt{p_i q_i}$$

Here  $p$  and  $q$  are eigenvalues of  $\rho_1$  and  $\rho_2$  respectively.

## Fidelity as function of eigenvalues

We start our analysis of fidelity by expressing it in terms of eigenvalues  $\lambda_i$ ,  $i = 1, \dots, N$  of the (positive) matrix  $\sqrt{\rho_1^{1/2} \rho_2 \rho_1^{1/2}}$ . Using the fact that matrix  $\rho_1 \rho_2$  is similar to matrix  $\sqrt{\rho_1} \rho_2 \sqrt{\rho_1}$  one can write

$$\sqrt{F(\rho_1, \rho_2)} = \text{tr} \sqrt{\sqrt{\rho_1} \rho_2 \sqrt{\rho_1}} = \sum_{i=1}^N \lambda_i,$$

and since  $\text{tr} \rho_1 \rho_2 = \text{tr} \sqrt{\rho_1} \rho_2 \sqrt{\rho_1} = \sum_{i=1}^N \lambda_i^2$  by squaring the above we get

$$F(\rho_1, \rho_2) = \left( \sum_{i=1}^N \lambda_i \right)^2 = \text{tr} \rho_1 \rho_2 + 2 \sum_{i < j} \lambda_i \lambda_j.$$

# Elementary symmetric functions

For a give matrix  $X \in \mathcal{M}_N$  with eigenvalues  $\lambda_1, \dots, \lambda_N$  we define elementary symmetric function  $s_m(X)$  as  $s_m(\lambda_1, \dots, \lambda_N)$

For example

$$s_2(X) = \sum_{i < j} \lambda_i \lambda_j,$$
$$s_3(X) = \sum_{i < j < k} \lambda_i \lambda_j \lambda_k.$$

Using this notion we can write the fidelity as

$$F(\rho_1, \rho_2) = \text{tr} \rho_1 \rho_2 + 2s_2(\sqrt{\sqrt{\rho_1} \rho_2 \sqrt{\rho_1}}).$$

## Lower bound by Uhlmann

In his unpublished work Uhlmann suggested an inequality

$$F(\rho_1, \rho_2) \geq \text{tr} \rho_1 \rho_2 + \sqrt{2} \sqrt{(\text{tr} \rho_1 \rho_2)^2 - \text{tr} \rho_1 \rho_2 \rho_1 \rho_2}.$$

We define sub-fidelity as

$$E(\rho_1, \rho_2) = \rho_1 \rho_2 + \sqrt{2} \sqrt{(\text{tr} \rho_1 \rho_2)^2 - \text{tr} \rho_1 \rho_2 \rho_1 \rho_2}.$$

Using elementary symmetric functions this quantity can be represented as

$$E(\rho_1, \rho_2) = \text{tr} \rho_1 \rho_2 + 2 \sqrt{s_2(\rho_1 \rho_2)}.$$



## Super-fidelity

We can introduce upper bound which is complementary to sub-fidelity<sup>1</sup>

$$F(\rho_1, \rho_2) \leq \text{tr}\rho_1\rho_2 + \sqrt{(1 - \text{tr}\rho_1^2)(1 - \text{tr}\rho_2^2)}.$$

Again we can use elementary symmetric function to get compact expression for super-fidelity

$$G(\rho_1, \rho_2) = \text{tr}\rho_1\rho_2 + \sqrt{(1 - \text{tr}\rho_1^2)(1 - \text{tr}\rho_2^2)} = \text{tr}\rho_1\rho_2 + 2\sqrt{s_2(\rho_1)s_2(\rho_2)}.$$

thus we have

$$s_2(\sqrt{\sqrt{\rho_1}\rho_2\sqrt{\rho_1}}) \leq \sqrt{s_2(\rho_1)s_2(\rho_2)}$$

<sup>1</sup>J. A. M, Z. Puchała, P. Horodecki, A. Uhlmann, K. Życzkowski, QI&C, **9**, 1&2 (2009), arXiv:0805.2037.

Two inequalities  $E(\rho_1, \rho_2) \leq F(\rho_1, \rho_2) \leq G(\rho_1, \rho_2)$  can be written in a compact way using elementary symmetric functions

$$\sqrt{s_2(\rho_1 \rho_2)} \leq s_2(\sqrt{\sqrt{\rho_1} \rho_2 \sqrt{\rho_1}}) \leq \sqrt{s_2(\rho_1) s_2(\rho_2)}$$

- ▶ If one of the states is pure we have equality  $E = F = G$ .
- ▶ Moreover these quantities coincide for one-qubit states ( $N = 2$ ).

## Properties of sub- and super-fidelity

Sub- and super-fidelity share some properties with fidelity

- i') **Bounds:**  $0 \leq E(\rho_1, \rho_2) \leq 1$  oraz  $0 \leq G(\rho_1, \rho_2) \leq 1$ .
- ii') **Symmetry:**  $E(\rho_1, \rho_2) = E(\rho_2, \rho_1)$  and  $G(\rho_1, \rho_2) = G(\rho_2, \rho_1)$ .
- iii') **Unitary invariance:**  $E(\rho_1, \rho_2) = E(U\rho_1 U^\dagger, U\rho_2 U^\dagger)$  and  $G(\rho_1, \rho_2) = G(U\rho_1 U^\dagger, U\rho_2 U^\dagger)$ , for any unitary  $U$ .
- iv') **Concavity:** Sub- and super-fidelity are concave,

$$E(A, \alpha B + (1 - \alpha)C) \geq \alpha E(A, B) + (1 - \alpha)E(A, C),$$

$$G(A, \alpha B + (1 - \alpha)C) \geq \alpha G(A, B) + (1 - \alpha)G(A, C).$$

- v') Super-fidelity (just like  $\sqrt{F}$ ) is **jointly concave** in its two arguments

$$\sqrt{F}(a\rho_1 + (1-a)\rho_2, a\rho'_1 + (1-a)\rho'_2) \geq a\sqrt{F}(\rho_1, \rho'_1) + (1-a)\sqrt{F}(\rho_2, \rho'_2)$$

for  $a \in [0, 1]$ .

# Properties of sub- and super-fidelity

## Properties for tensor products

Both  $E$  and  $G$  are not multiplicative. On the other hand

# Properties of sub- and super-fidelity

## Properties for tensor products

Both  $E$  and  $G$  are not multiplicative. On the other hand  
**vi')** Super-fidelity is super-multiplicative:

$$G(\rho_1 \otimes \rho_2, \rho_3 \otimes \rho_4) \geq G(\rho_1, \rho_3)G(\rho_2, \rho_4),$$

# Properties of sub- and super-fidelity

## Properties for tensor products

Both  $E$  and  $G$  are not multiplicative. On the other hand  
**vi')** Super-fidelity is super-multiplicative:

$$G(\rho_1 \otimes \rho_2, \rho_3 \otimes \rho_4) \geq G(\rho_1, \rho_3)G(\rho_2, \rho_4),$$

**vii')** Sub-fidelity is sub-multiplicative

$$E(\rho_1 \otimes \rho_2, \rho_3 \otimes \rho_4) \leq E(\rho_1, \rho_3)E(\rho_2, \rho_4).$$

## Classical case

If the density matrices  $\rho_p$  and  $\rho_q$  commute, discussed bound can be expressed in terms of respective eigenvalues  $\{p_i\}_{i=1}^N$  and  $\{q_i\}_{i=1}^N$ :

$$E(\rho_p, \rho_q) = \sum_{i=1}^N p_i q_i + \sqrt{2 \left[ \left( \sum_{i=1}^N p_i q_i \right)^2 - \sum_{i=1}^N p_i^2 q_i^2 \right]},$$

$$F(\rho_p, \rho_q) = \left( \sum_{i=1}^N \sqrt{p_i q_i} \right)^2,$$

$$G(\rho_p, \rho_q) = \sum_{i=1}^N p_i q_i + \sqrt{\left( 1 - \sum_{i=1}^N p_i^2 \right) \left( 1 - \sum_{i=1}^N q_i^2 \right)}.$$

## Mixed states

To get some feeling about behaviour of  $E$  and  $G$  we calculate them for states of the form

$$\rho_a = a|\psi\rangle\langle\psi| + (1-a)I/N.$$

where  $|\psi\rangle$  is an arbitrary pure state.

For  $\rho_* := I/N$  we get

$$F(\rho_a, \rho_*) = \frac{1}{N^2} \left( \sqrt{(N-1)a+1} + (N-1)\sqrt{1-a} \right)^2,$$





and sub- and super-fidelity are expressed as

$$E(\rho_a, \rho_*) = \frac{1}{N} + \sqrt{2} \frac{1}{N} \sqrt{1 - \frac{1}{N}} \sqrt{1 - a^2},$$

$$G(\rho_a, \rho_*) = \frac{1}{N} + \left(1 - \frac{1}{N}\right) \sqrt{1 - a^2}.$$



## Comparison of sub-fidelity $E$ , fidelity $F$ and super-fidelity $G$ .



## Difference $G - F$ and $E - F$

$F$  and  $G$  coincide if one of the states is pure, but it is natural to ask how big the difference  $G - F$  might be.

Let us use the Hilbert space of dimension  $N = 2M$  and states  $\rho_1 = \frac{2}{N} \text{diag}(1, \dots, 1, 0, \dots, 0)$  and  $\rho_2 = \frac{2}{N} \text{diag}(0, \dots, 0, 1, \dots, 1)$ .

## Difference $G - F$ and $E - F$


$F$  and  $G$  coincide if one of the states is pure, but it is natural to ask how big the difference  $G - F$  might be.

Let us use the Hilbert space of dimension  $N = 2M$  and states  $\rho_1 = \frac{2}{N} \text{diag}(1, \dots, 1, 0, \dots, 0)$  and  $\rho_2 = \frac{2}{N} \text{diag}(0, \dots, 0, 1, \dots, 1)$ . Since they are supported by orthogonal subspaces their fidelity vanishes,  $F(\rho_1, \rho_2) = 0$ . On the other hand their super-fidelity is equal to

$$G(\rho_1, \rho_2) = \frac{N - 2}{N},$$

and the difference  $F - G$  can be arbitrarily close to 1.


# Maximal difference



Max difference between fidelity and sub- and super- fidelity for random states in dimensions  $N = 2, 3, \dots, 62$ .

# Average difference

On average situation looks somehow better.



Average difference between fidelity and sub- and super- fidelity for some values of  $N \in [2, 62]$ .

## Super-fidelity and trace distance

For any  $\rho_1, \rho_2 \in \Omega_N$  super-fidelity and trace distance are related by the inequality<sup>2</sup>

$$1 - G(\rho_1, \rho_2) \leq D_{\text{tr}}(\rho_1, \rho_2)$$

Probability of error for distinguishing two density matrices  $\rho_1, \rho_2 \in \Omega_N$  is expressed by the trace distance as

$$P_E(\rho_1, \rho_2) = \frac{1}{2}(1 - D_{\text{tr}}(\rho_1, \rho_2)).$$

Using the above inequalities we can write

$$\frac{1}{2}G(\rho_1, \rho_2) \geq P_E(\rho_1, \rho_2)$$

---

<sup>2</sup>Z. Puchała, J. A. M., *Bounds on trace distance based on super-fidelity, in preparation*

## Experimental setup for measuring super-fidelity

We use fact that  $\text{tr} V_{12} \rho_1 \otimes \rho_2 = \text{tr} \rho_1 \rho_2$  where  $V_{12}$  is a SWAP operator.  $V_{12} = P_{12}^+ - P_{12}^-$  is hermitian and thus represents an observable.


To measure  $G$  we need a source which creates pairs  $\rho_i \otimes \rho_j$ ,  $i, j = 1, 2$ .

The probability of measuring  $P_{12}^-$  reads  $p_{ij}^- = \text{tr} P_{12}^- \rho_i \otimes \rho_j$  and using it we can write

$$G = 1 - 2(p_{12}^- - \sqrt{p_{11}^- - p_{22}^-})$$

## Experimental setup for measuring super-fidelity

Probability of the event that both detectors click is equal to  $p_{ij}^-$ .  
 On detectors clicks with  $p_{ij}^+ = 1 - p_{ij}^-$ . Beam-splitter projects on  $P^-$  or  $P^+$




The experimental setup is in this case very simple.<sup>3</sup>

<sup>3</sup>F. A. Bovino et al, PRL **95**, 240407 (2005)



# Computational efficiency

$E$  and  $G$  are much easier to calculate than fidelity  $F$ . To compute any of these bounds it is enough to evaluate three traces only.






(See also P. E. M. F. Mendonca, *et al*, arXiv:0806.1150)

# Conclusions

- ▶ Proposed bounds share with fidelity its main features (they are bounded, symmetric, unitary invariant and concave).
- ▶ Super-fidelity  $G$  can be used in place of fidelity  $F$  for small systems or when at least one of the states is pure enough.
- ▶ Sub- and super-fidelity can be (in principle) measured in laboratory.
- ▶ It is easy to calculate them using standard computer algebra systems.

## References

-  J. A. M, Z. Puchała, P. Horodecki, A. Uhlmann, K. Życzkowski, *Sub- and super-fidelity as bounds for quantum fidelity*, Quantum Information & Computation, **9**, No.1&2 (2009), arXiv:0805.2037.
-  P. E. M. F. Mendonca, R. d. J. Napolitano, M. A. Marchioli, C. J. Foster, and Y.-C. Liang, *An alternative fidelity measure for quantum states*, to appear in PRA, arXiv:0806.1150
-  J.-L. Chen, L. Fu, A.A. Ungar, and X.-G. Zhao, *Alternative fidelity measure between two states of an  $N$ -state quantum system*, *Phys. Rev. A* **65**, 054304 (2002).

Thank you.